


12I – Parallelism

If two vectors are parallel, then one is a scalar multiple of the other.

Given any non-zero vector \mathbf{v} and non-zero scalar k , the vector $k\mathbf{v}$ is parallel to \mathbf{v} .



- If \mathbf{a} is parallel to \mathbf{b} , then there exists a scalar k such that $\mathbf{a} = k\mathbf{b}$.
- If $\mathbf{a} = k\mathbf{b}$ for some scalar k , then
 - ▶ \mathbf{a} is parallel to \mathbf{b} , and
 - ▶ $|\mathbf{a}| = |k| |\mathbf{b}|$.

Example:

1. If $\mathbf{a} = \begin{pmatrix} 15 \\ 10 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$ then \mathbf{a} is parallel to \mathbf{b} .

$$|\vec{a}| = 5|\vec{b}|$$

Find: $|\mathbf{a}|$ and $|\mathbf{b}|$

$$\begin{aligned} |\vec{a}| &= \sqrt{(15)^2 + (10)^2} \\ &= \sqrt{225 + 100} \\ &= \sqrt{325} \\ &= 5\sqrt{13} \end{aligned}$$

$$\begin{aligned} |\vec{b}| &= \sqrt{(3)^2 + (2)^2} \\ &= \sqrt{9 + 4} \\ &= \sqrt{13} \end{aligned}$$

2. Find r and s given that $\mathbf{a} = \begin{pmatrix} 2 \\ -1 \\ r \end{pmatrix}$ is parallel to $\mathbf{b} = \begin{pmatrix} 8 \\ s \\ -6 \end{pmatrix}$.

$$k\vec{\mathbf{a}} = \vec{\mathbf{b}}$$

$$k \begin{pmatrix} 2 \\ -1 \\ r \end{pmatrix} = \begin{pmatrix} 8 \\ s \\ -6 \end{pmatrix}$$

$$2k = 8$$
$$k = 4$$

$$-k = s$$

$$-4 = s$$

$$kr = -6$$

$$4r = -6$$

$$r = \frac{-6}{4} = -\frac{3}{2}$$

Unit Vectors:

Given a non-zero vector \mathbf{v} , its magnitude $|\mathbf{v}|$ is a scalar quantity.

If we multiply \mathbf{v} by the scalar $\frac{1}{|\mathbf{v}|}$, we obtain the parallel vector $\frac{1}{|\mathbf{v}|} \mathbf{v}$.

The length of this vector is $\left| \frac{1}{|\mathbf{v}|} \right| |\mathbf{v}| = \frac{|\mathbf{v}|}{|\mathbf{v}|} = 1$, so $\frac{1}{|\mathbf{v}|} \mathbf{v}$ is a unit vector in the direction of \mathbf{v} .

- A unit vector in the direction of \mathbf{v} is $\frac{1}{|\mathbf{v}|} \mathbf{v}$.
- A vector \mathbf{b} of length k in the same direction as \mathbf{a} is $\mathbf{b} = \frac{k}{|\mathbf{a}|} \mathbf{a}$.
- A vector \mathbf{b} of length k which is *parallel to* \mathbf{a} could be $\mathbf{b} = \pm \frac{k}{|\mathbf{a}|} \mathbf{a}$.

6 Find the unit vector in the direction of:

a $\mathbf{i} + 2\mathbf{j}$

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$|\vec{i} + 2\vec{j}| = \sqrt{(1)^2 + (2)^2}$$

$$= \sqrt{5}$$

$$\text{Unit vector} = \frac{1}{\sqrt{5}} (\vec{i} + 2\vec{j})$$

b ~~$2\mathbf{i} - 3\mathbf{k}$~~

$$\begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix}$$

c $2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$

$$|2\vec{i} - 2\vec{j} + \vec{k}| = \sqrt{(2)^2 + (-2)^2 + (1)^2}$$

$$= \sqrt{4 + 4 + 1}$$

$$= \sqrt{9}$$

$$= 3$$

$$\text{Unit vector} = \frac{1}{3} (2\vec{i} - 2\vec{j} + \vec{k})$$

10 Find a vector \mathbf{b} in:

a the same direction as $\sqrt{2} \begin{pmatrix} -1 \\ 4 \\ 1 \end{pmatrix}$ and with length 6 units

① find $|\vec{v}|$

$$|\vec{v}| = \sqrt{(-1)^2 + (4)^2 + (1)^2}$$

$$= \sqrt{1+16+1}$$

$$= \sqrt{18}$$

$$= 3\sqrt{2}$$

② unit vector in the same direction is $\frac{1}{3\sqrt{2}} \begin{pmatrix} -1 \\ 4 \\ 1 \end{pmatrix}$

③ Of length 6 will be 6 times the unit vector

$$\vec{b} = \frac{6}{3\sqrt{2}} \begin{pmatrix} -1 \\ 4 \\ 1 \end{pmatrix}$$

$$= \frac{2}{\sqrt{2}} \begin{pmatrix} -1 \\ 4 \\ 1 \end{pmatrix} = \sqrt{2} \begin{pmatrix} -1 \\ 4 \\ 1 \end{pmatrix}$$

12J – The Scalar Product of Two Vectors



- ▶ The scalar product of 2 vectors, which results in a scalar answer and has the notation $\mathbf{v} \bullet \mathbf{w}$ (read “v dot w”).
- ▶ The vector product of 2 vectors, which results in a vector answer and has the notation $\mathbf{v} \times \mathbf{w}$ (read “v cross w”).

The scalar product of two vectors is also known as the dot product or inner product.


If $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$, the scalar product of \mathbf{v} and \mathbf{w} is defined as

$$\mathbf{v} \bullet \mathbf{w} = v_1 w_1 + v_2 w_2 + v_3 w_3.$$

Angle Between Vectors:

Consider the vectors  $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ and  $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$.

We translate one of the vectors so that they both originate from the same point.



This vector is $-\mathbf{v} + \mathbf{w} = \mathbf{w} - \mathbf{v}$
and has length $|\mathbf{w} - \mathbf{v}|$.

Using the cosine rule, $|\mathbf{w} - \mathbf{v}|^2 = |\mathbf{v}|^2 + |\mathbf{w}|^2 - 2|\mathbf{v}||\mathbf{w}|\cos\theta$

$$\text{But } \mathbf{w} - \mathbf{v} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} - \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} w_1 - v_1 \\ w_2 - v_2 \\ w_3 - v_3 \end{pmatrix}$$

$$\therefore (w_1 - v_1)^2 + (w_2 - v_2)^2 + (w_3 - v_3)^2 = v_1^2 + v_2^2 + v_3^2 + w_1^2 + w_2^2 + w_3^2 - 2|\mathbf{v}||\mathbf{w}|\cos\theta$$

$$\therefore \underline{v_1 w_1 + v_2 w_2 + v_3 w_3} = |\mathbf{v}||\mathbf{w}|\cos\theta$$

$$\therefore \underline{\mathbf{v} \bullet \mathbf{w}} = |\mathbf{v}||\mathbf{w}|\cos\theta$$

The angle θ between two vectors \mathbf{v} and \mathbf{w} can be found using

$$\cos \theta = \frac{\mathbf{v} \bullet \mathbf{w}}{|\mathbf{v}| |\mathbf{w}|}$$

ALGEBRAIC PROPERTIES OF THE SCALAR PRODUCT

The scalar product has the following algebraic properties for both 2-D and 3-D vectors:

▶ $\mathbf{v} \bullet \mathbf{w} = \mathbf{w} \bullet \mathbf{v}$



▶ $\mathbf{v} \bullet \mathbf{v} = |\mathbf{v}|^2$

▶ $\mathbf{v} \bullet (\mathbf{w} + \mathbf{x}) = \mathbf{v} \bullet \mathbf{w} + \mathbf{v} \bullet \mathbf{x}$

▶ $(\mathbf{v} + \mathbf{w}) \bullet (\mathbf{x} + \mathbf{y}) = \mathbf{v} \bullet \mathbf{x} + \mathbf{v} \bullet \mathbf{y} + \mathbf{w} \bullet \mathbf{x} + \mathbf{w} \bullet \mathbf{y}$

GEOMETRIC PROPERTIES OF THE SCALAR PRODUCT

- ▶ For non-zero vectors \mathbf{v} and \mathbf{w} :
 $\mathbf{v} \bullet \mathbf{w} = 0 \iff \mathbf{v}$ and \mathbf{w} are perpendicular or orthogonal.
- ▶ $|\mathbf{v} \bullet \mathbf{w}| = |\mathbf{v}| |\mathbf{w}| \iff \mathbf{v}$ and \mathbf{w} are non-zero parallel vectors.
- ▶ If θ is the angle between vectors \mathbf{v} and \mathbf{w} then: $\mathbf{v} \bullet \mathbf{w} = |\mathbf{v}| |\mathbf{w}| \cos \theta$
If θ is acute, $\cos \theta > 0$ and so $\mathbf{v} \bullet \mathbf{w} > 0$
If θ is obtuse, $\cos \theta < 0$ and so $\mathbf{v} \bullet \mathbf{w} < 0$.
The angle between two vectors is always taken as the angle θ such that $0^\circ \leq \theta \leq 180^\circ$, rather than reflex angle α .



The first two of these results can be demonstrated as follows:

If \mathbf{v} is perpendicular to \mathbf{w} then $\theta = 90^\circ$.

If \mathbf{v} is parallel to \mathbf{w} then $\theta = 0^\circ$ or 180° .

$$\begin{aligned}\therefore \mathbf{v} \bullet \mathbf{w} &= |\mathbf{v}| |\mathbf{w}| \cos \theta \\ &= |\mathbf{v}| |\mathbf{w}| \cos 90^\circ \\ &= 0\end{aligned}$$

$$\begin{aligned}\therefore \mathbf{v} \bullet \mathbf{w} &= |\mathbf{v}| |\mathbf{w}| \cos \theta \\ &= |\mathbf{v}| |\mathbf{w}| \cos 0^\circ \text{ or } |\mathbf{v}| |\mathbf{w}| \cos 180^\circ \\ &= \pm |\mathbf{v}| |\mathbf{w}| \\ \therefore |\mathbf{v} \bullet \mathbf{w}| &= |\mathbf{v}| |\mathbf{w}|\end{aligned}$$


To formally prove these results we must also show that their converses are true.

Examples:

1. Find the scalar product of $\vec{p} = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ and $\vec{q} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$.

$$\begin{aligned}\vec{p} \cdot \vec{q} &= \begin{pmatrix} 1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ 5 \end{pmatrix} \\ &= (1)(-2) + (3)(5) \\ &= -2 + 15 \\ &= 13\end{aligned}$$

2. Find m such that $\vec{p} = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$ and $\vec{q} = \begin{pmatrix} 1 \\ m \end{pmatrix}$ are perpendicular.



$$\vec{p} \cdot \vec{q} = 0$$

$$\begin{pmatrix} -2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ m \end{pmatrix} = 0$$


$$(-2)(1) + (3)(m) = 0$$

$$-2 + 3m = 0$$

$$3m = 2$$

$$m = \frac{2}{3}$$

3. Find the form of all vectors which are perpendicular to $\begin{pmatrix} 3 \\ 7 \end{pmatrix} = \vec{a}$



$$\begin{pmatrix} 3 \\ 7 \end{pmatrix} \cdot \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = 0$$

$$3b_1 + 7b_2 = 0$$

$$\begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \begin{pmatrix} -7 \\ 3 \end{pmatrix}$$

$$\begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \begin{pmatrix} 7 \\ -3 \end{pmatrix}$$

$\vec{b} = \begin{pmatrix} -7 \\ 3 \end{pmatrix}$ is perpendicular to $\begin{pmatrix} 3 \\ 7 \end{pmatrix}$

all vectors would be $k \begin{pmatrix} -7 \\ 3 \end{pmatrix}$ $k \neq 0$

$m \begin{pmatrix} 7 \\ -3 \end{pmatrix}$ $m \neq 0$

4. Find $\mathbf{a} \cdot \mathbf{b}$ for:

(a) $|\mathbf{a}| = 4$, $|\mathbf{b}| = 5$, and $\theta = 150^\circ$

$$\begin{aligned}\vec{a} \cdot \vec{b} &= (4)(5) \cos 150^\circ \\ &= 20 \cos 150^\circ \\ &= 20 \left(-\frac{\sqrt{3}}{2} \right) \\ &= -10\sqrt{3}\end{aligned}$$

(b) $|\mathbf{a}| = 3$, $|\mathbf{b}| = 2$, and $\theta = 75^\circ$

$$\begin{aligned}\vec{a} \cdot \vec{b} &= (3)(2) \cos 75^\circ \\ &= 6 \cos 75^\circ \\ &\approx 1.5529\end{aligned}$$

$$\vec{a} \cdot \vec{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$$



5. Find the angle between $\vec{a} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ and $\vec{b} = \begin{pmatrix} 2 \\ 7 \end{pmatrix}$.

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

$$\cos \theta = \frac{(4)(2) + (-3)(7)}{\sqrt{(4)^2 + (-3)^2} \sqrt{(2)^2 + (7)^2}}$$

$$= \frac{8 - 21}{(\sqrt{25})(\sqrt{53})}$$


$$\cos \theta = \frac{-13}{5\sqrt{53}}$$

$$\cos \theta = -0.857137$$

$$\theta = 110.924^\circ$$

$$\theta = 110.92^\circ$$

6. Find the measure of angle ABC for A(2, -1), B(3, 4) and C(-1, 3).



$$\vec{BA} \cdot \vec{BC} = |\vec{BA}| |\vec{BC}| \cos \theta$$

$$\cos \theta = \frac{\vec{BA} \cdot \vec{BC}}{|\vec{BA}| |\vec{BC}|}$$

$$= \frac{(-1)(-4) + (5)(-1)}{(\sqrt{(-1)^2 + (5)^2})(\sqrt{(-4)^2 + (-1)^2})}$$

Find \vec{BA} , \vec{BC}

$$\begin{aligned} \vec{BA} &= \begin{pmatrix} 2-3 \\ -1-4 \end{pmatrix} \\ &= \begin{pmatrix} -1 \\ -5 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \vec{BC} &= \begin{pmatrix} -1-3 \\ 3-4 \end{pmatrix} \\ &= \begin{pmatrix} -4 \\ -1 \end{pmatrix} \end{aligned}$$

$$\cos \theta = \frac{+9}{\sqrt{26} \cdot \sqrt{17}}$$

$$\theta = 64.65^\circ$$